

THE ALEXANDER POLYNOMIAL OF $(1, 1)$ -KNOTS

A. CATTABRIGA

*Mathematics Department, University of Bologna,
P.zza di Porta S. Donato, 5, 40126 Bologna, Italy
cattabri@dm.unibo.it*

ABSTRACT

In this paper we investigate the Alexander polynomial of $(1, 1)$ -knots, which are knots lying in a 3-manifold with genus one at most, admitting a particular decomposition. More precisely, we study the connections between the Alexander polynomial and a polynomial associated to a cyclic presentation of the fundamental group of an n -fold strongly-cyclic covering branched over the knot K , which we call the n -cyclic polynomial of K . In this way, we generalize to all $(1, 1)$ -knots, with the only exception of those lying in $\mathbf{S}^2 \times \mathbf{S}^1$, a result obtained by J. Minkus for 2-bridge knots and extended by the author and M. Mulazzani to the case of $(1, 1)$ -knots in $\mathbf{S}^3$. As corollaries some properties of the Alexander polynomial of knots in $\mathbf{S}^3$ are extended to the case of $(1, 1)$ -knots in lens spaces.

Keywords: $(1, 1)$ -knots, cyclic branched coverings, Alexander polynomial, cyclically presented groups.

Mathematics Subject Classification 2000: Primary 57M12, 57M27; Secondary 57N10.

1. Introduction

A knot K in a 3-manifold M is called a $(1, 1)$ -knot, or a 1-bridge torus knot, if it can be embedded into a Heegaard torus of M except at one over (or under) bridge. This means that there exists a Heegaard splitting of genus one

$$(M, K) = (H, A) \cup_{\varphi} (H', A'), \quad (1.1)$$

where H and H' are solid tori, $A \subset H$ and $A' \subset H'$ are properly embedded trivial arcs¹, and $\varphi : (\partial H', \partial A') \rightarrow (\partial H, \partial A)$ is an attaching homeomorphism (see Fig. 1.1). Such a decomposition is called $(1, 1)$ -*decomposition* of the knot. By definition, the ambient manifold has genus less or equal than one, so M , up to homeomorphism,

¹This means that there exists a disk $D \subset H$ (resp. $D' \subset H'$) with $A \cap D = A \cap \partial D = A$ and $\partial D - A \subset \partial H$ (resp. $A' \cap D' = A' \cap \partial D' = A'$ and $\partial D' - A' \subset \partial H'$).

is $\mathbf{S}^2 \times \mathbf{S}^1$ or a lens space $L(p, q)$, including the case $L(1, 0) = \mathbf{S}^3$. Important examples of $(1, 1)$ -knots in $\mathbf{S}^3$ are torus knots (by definition) and 2-bridge knots (see [15]). This family of knots has interesting features and has recently been studied by many authors from different points of view. For instance, in [5, 7, 9], different representations of $(1, 1)$ -knots are given, while in [1, 4, 6, 13, 17] their connections with manifolds with cyclically presented fundamental groups are studied.




Fig. 1.1. A $(1, 1)$ -decomposition.

In this paper we investigate the Alexander polynomial of $(1, 1)$ -knots. This class of knots seems to have very interesting features with regards to the Alexander polynomial. For example, in [12], the author shows that it is possible to realize each Alexander polynomial in $\mathbf{S}^3$ using $(1, 1)$ -knots. Moreover, he exhibits an infinite family of $(1, 1)$ -knots in $\mathbf{S}^3$ with a trivial Alexander polynomial. In this paper we focus especially on $(1, 1)$ -knots in lens spaces. From the results of [17], each n -fold strongly-cyclic branched covering of a $(1, 1)$ -knot admits a cyclic presentation for the fundamental group and a polynomial associated to this presentation. Moreover, if $K \subset L(p, q)$ and n is coprime with p , the covering is unique, and so there is a natural way to associate to K a polynomial that will be called the n -cyclic polynomial of K . Using the reduced Reidemeister torsion of the knot complement, we find the relation between this polynomial and the Alexander polynomial. In this way, we generalize to all $(1, 1)$ -knots, with the only exception of those lying in $\mathbf{S}^2 \times \mathbf{S}^1$, a result of [16] for 2-bridge knots and of [4] for $(1, 1)$ -knots in $\mathbf{S}^3$. As corollaries, we extend to all $(1, 1)$ -knots in lens spaces some properties of the Alexander polynomial of knots in $\mathbf{S}^3$.

In Section 2 we give the definition of n -cyclic polynomials. In Section 3 we recall the definition of the Alexander polynomial of a knot, and some results concerned with it. The main result of this paper is established in Section 4, where some corollaries and examples are also collected.

2. Strongly-cyclic coverings and cyclic presentations of groups

Let F_n denote the free group of rank n . A group G is called *cyclically presented* if there exists $n > 0$ and $w \in F_n$ such that

$$G = G_n(w) = \langle x_1, \dots, x_n \mid w, \theta_n(w), \dots, \theta_n^{n-1}(w) \rangle, \quad (2.1)$$

where $\theta_n : F_n \rightarrow F_n$ is the automorphism defined by $\theta_n(x_i) = x_{i+1}$ (subscripts mod n), for $i = 1, \dots, n$. Such a presentation is called a *cyclic presentation*. Obviously $G_n(w)$ has an automorphism of order n induced by θ_n .

The *polynomial associated to the cyclic presentation* $G_n(w)$ is

$$p_{n,w}(t) = \sum_{i=1}^n a_i t^{i-1}, \quad (2.2)$$

where a_i is the total exponent sum of the letter x_i in the word w , for $i = 1, \dots, n$.

For each $s = 0, \dots, (n-1)$, we have $G_n(w) = G_n(\theta_n^s(w))$, while the polynomials $p_{n,\theta_n^s(w)}(t)$ are different elements of $\mathbb{Z}[t]$. However, their image in the quotient ring $\mathbb{Z}[t]/(t^n - 1)$ is the same, up to units. So, from now on, we will consider the polynomial associated to the cyclic presentation $G_n(w)$ as an element of $\mathbb{Z}[t]/(t^n - 1)$.

Note that the abelianization of $G_n(w)$ has a circulant presentation matrix (as a $\mathbb{Z}$ -module), whose first row is given by the coefficients of $p_{n,w}$.

An n -fold cyclic covering $f : \widetilde{M} \rightarrow M$, branched over a knot $K \subset M$, is called *strongly-cyclic* if the branched index of K is n . In other words², if $\omega_f : H_1(M - N(K)) \rightarrow \mathbb{Z}_n$ denotes the monodromy of f and $\overline{m}$ is the first homology class of a meridian loop of K , $\omega_f(\overline{m})$ generates $\mathbb{Z}_n$. Moreover, two n -fold strongly-cyclic coverings f and f' , branched over the same knot $K \subset M$, are equivalent if there exists an invertible element $u \in \mathbb{Z}_n$ such that the following diagram commutes

$$\begin{array}{ccc} H_1(L(p, q) - N(K)) & \xrightarrow{\omega_{f'}} & \mathbb{Z}_n \\ \omega_f \downarrow & \nearrow \mu_u & \\ \mathbb{Z}_n & & \end{array} \quad (2.3)$$

where μ_u denotes the multiplication by u . So, up to equivalence, a cyclic branched covering f is strongly-cyclic if $\omega_f(\overline{m}) = 1$. From now on, $C_n(K)$ will denote an n -fold strongly-cyclic branched covering of K .

If we are dealing with knots in $\mathbf{S}^3$, every n -fold cyclic branched covering of a knot is strongly-cyclic, so it exists and is unique, up to equivalence. This is no longer true for a $(1, 1)$ -knot K in a lens space $L(p, q)$. However, from the existence

²We denote with $N(K) \subset M$ an open tubular neighborhood of K in M .

and uniqueness conditions of [4], it follows that for each n coprime with p there is a unique n -fold strongly-cyclic branched covering of K .

The connection between $(1, 1)$ -knots and cyclically presented groups is given by the following theorem.

Theorem 2.1. [17] *Every n -fold strongly-cyclic branched covering of a $(1, 1)$ -knot admits a cyclic presentation for the fundamental group with n generators.*

Moreover, this presentation is obtained by lifting a $(1, 1)$ -presentation of the knot, as follows (for details see [4]).

From a $(1, 1)$ -decomposition $(L(p, q), K) = (H, A) \cup_{\varphi} (H', A')$ of K , we get the following presentation of the knot group:

$$\pi_1(L(p, q) - N(K), *) = \langle \alpha, \gamma, \mid r(\alpha, \gamma) \rangle, \quad (2.4)$$

where the generators α and γ represent, respectively, a longitude of ∂H and a meridian loop of K , and the relator $r(\alpha, \gamma)$ corresponds to the loop $\varphi(\partial D')$, where D' is a meridian disk of H' that does not intersect A' (see Fig. 1.1).

If necessary, by replacing the covering $f : \widetilde{M} \rightarrow L(p, q)$ with an equivalent one, we can suppose that $\omega_f(\gamma) = 1$. Let $\bar{r}(x, \gamma) = r(x\gamma^{\omega_f(\alpha)}, \gamma) = x^{\varepsilon_1}\gamma^{\delta_1} \dots x^{\varepsilon_s}\gamma^{\delta_s}$ for some $\varepsilon_1, \dots, \varepsilon_s, \delta_1, \dots, \delta_s \in \mathbb{Z}$. We have $\pi_1(\widetilde{M}, *) = G_n(w)$ where:

$$w = x_{i_1}^{\varepsilon_1} \dots x_{i_s}^{\varepsilon_s} \quad (\text{subscripts mod } n), \quad (2.5)$$

with $i_k \equiv 1 + \sum_{j=1}^{k-1} \delta_j \pmod{n}$, for $k = 1, \dots, s$.

For each n coprime with p , the polynomial associated to the cyclic presentation of the (unique) n -fold strongly-cyclic branched covering of K obtained as above (i.e. lifting a $(1, 1)$ -decomposition of K and under the condition $\omega_f(\gamma) = 1$) will be called the n -cyclic polynomial of K , and it will be denoted with $\Gamma_{K,n}$.

Remark 2.1. In the case of a $(1, 1)$ -knot $K \subset \mathbf{S}^2 \times \mathbf{S}^1$, which corresponds to $p = 0$, the situation is rather different. We do not have the uniqueness of the n -fold strongly-cyclic branched covering of K for any value of $n > 1$, and so we do not have a natural way of defining the n -cyclic polynomial. Moreover, there exists at most a finite number of finite strongly-cyclic branched covering of K , for almost all the K .

Let $\Delta_K \in \mathbb{Z}[t, t^{-1}]$ be the Alexander polynomial of a knot $K \subset \mathbf{S}^3$ and denote with $\Delta_{K,n}$ its projection on $\mathbb{Z}[t]/(t^n - 1)$. With these notations the results of [4] and [16] can be restated as follows.

Proposition 2.1. *Let $K \subset \mathbf{S}^3$. The following holds:*

- (i) [16] *if K is a 2-bridge knot, for each $n > 1$, we have $\Gamma_{K,n}(t) = \Delta_{K,n}(t)$, up to units of $\mathbb{Z}[t]/(t^n - 1)$.*
- (ii) [4] *if K is a $(1, 1)$ -knot, for each $n > 1$, we have $\Gamma_{K,n}(t) = \Delta_{K,n}(t)$, up to units of $\mathbb{Z}[t]/(t^n - 1)$.*

The main result of this article is the generalization of this relation to all $(1, 1)$ -knots in lens spaces.

3. Alexander polynomial

In this section we recall the definition of the Alexander polynomial of a knot K in a compact, connected 3-manifold, and give some of its characteristics.

Let M be a compact connected manifold, $*$ $\in M$ be a fixed point and denote with $\Phi : \pi_1(M, *) \rightarrow H_1(M)$ the Hurewitz homomorphism. Consider the projection $j : H_1(M) \rightarrow H_1(M)/\text{Tors}(H_1(M))$ and the induced ring homomorphism³ $\bar{j} : \mathbb{Z}[H_1(M)] \rightarrow \mathbb{Z}[H_1(M)/\text{Tors}(H_1(M))]$. Note that, if r is the first Betti number of M , we have $\mathbb{Z}[H_1(M)/\text{Tors}(H_1(M))] \cong \mathbb{Z}[t_1, t_1^{-1}, \dots, t_r, t_r^{-1}]$, where $t_1, \dots, t_r$ are generators of $H_1(M)/\text{Tors}(H_1(M))$. Let $E_1(M) \subset \mathbb{Z}[H_1(M)]$ be the first elementary ideal of $\pi_1(M, *)$ (see [10]) and denote with $\bar{E}_1(M)$ the smallest principal ideal of $\mathbb{Z}[H_1(M)/\text{Tors}(H_1(M))]$ containing $\bar{j}(E_1(M))$. The generator Δ_M of $\bar{E}_1(M)$ is well-defined up to multiplication by units of $\mathbb{Z}[H_1(M)/\text{Tors}(H_1(M))]$ and is called the *Alexander polynomial* of M .

If $K \subset X$ is a knot in a compact connected 3-manifold, the Alexander polynomial of K is the Alexander polynomial of $M = X - N(K)$.

For a knot $K \subset \mathbf{S}^3$ the Alexander polynomial has the following characterization.

Theorem 3.1. [10] *Let $\Delta_K(t) \in \mathbb{Z}[t, t^{-1}]$ be the Alexander polynomial of a knot $K \subset \mathbf{S}^3$. We have:*

- (i) $\Delta_K(t) = \Delta_K(t^{-1})$.
- (ii) $\Delta_K(1) = \pm 1$.

Moreover, each polynomial $q(t) \in \mathbb{Z}[t, t^{-1}]$ satisfying conditions 1 and 2 is the Alexander polynomial of a knot in $\mathbf{S}^3$.

Actually, from [12], each Alexander polynomial of a knot in $\mathbf{S}^3$ can be realized by a $(1, 1)$ -knot in $\mathbf{S}^3$.

Moreover, the Alexander polynomial of a knot $K \subset \mathbf{S}^3$ determines the order of the first homology group of $C_n(K)$, the n -fold (strongly-)cyclic branched covering of K .

Theorem 3.2. [11] *The abelian group $H_1(C_n(K))$ is finite if and only if no root of the Alexander polynomial $\Delta_K(t)$ of K is an n -root of unity. In this case:*

$$\sharp H_1(C_n(K)) = \left| \prod_{\zeta^n=1} \Delta_K(\zeta) \right|. \quad (3.1)$$

Let $K \subset L(p, q)$ be a $(1, 1)$ -knot. Then $H_1(L(p, q) - N(K)) \cong \mathbb{Z} \oplus \mathbb{Z}_d$, where $d > 0$ divides p (see [4]). Then the first Betti number of $L(p, q) - N(K)$ is one, so Δ_K lies in $\mathbb{Z}[t, t^{-1}]$. Moreover, it is easy to check that $\chi(L(p, q) - N(K)) = 0$. So the following theorem holds with $M = L(p, q) - N(K)$.

³For a group G we denote with $\mathbb{Z}[G]$ its integral group ring.

Theorem 3.3. [20, Theorem B] *Let M be a compact, connected, orientable 3-manifold with $\chi(M) = 0$. If $\partial M \neq \emptyset$ and $b_1(M) = 1$, then $r_M(t) = \Delta_M(t)/(t-1)$, where $r_M(t) \in \mathbb{Q}(t)$ denotes the reduced Reidemeister torsion of M and t is a generator of $H_1(M)/\text{Tors}(H_1(M))$.*

From this theorem it follows (see [20, 21]) that for the Alexander polynomial of $(1,1)$ -knots in lens spaces it is still true that $\Delta_K(t) = \Delta_K(t^{-1})$. However, we will see that the property $\Delta_K(1) = \pm 1$ is no longer true.

For details on the reduced Reidemeister torsion, and on its calculation, we refer to [20, 21].

Remark 3.1. In the case of a $(1,1)$ -knot $K \subset \mathbf{S}^2 \times \mathbf{S}^1$ it could happen that the first Betti number is two. That is the case, for example, of the trivial knot.

4. Main theorem

Theorem 4.1. *Let $K \subset L(p, q)$ be a $(1,1)$ -knot and denote with d the order of the torsion subgroup of $H_1(L(p, q) - N(K))$. Then, for each $n > 1$ such that $\gcd(p, n) = 1$, we have:*

$$\Gamma_{K,n}(t^{\frac{p}{d}}) = \Delta_{K,n}(t) \sum_{i=0}^{\frac{p}{d}-1} t^i, \quad (4.1)$$

up to units of $\mathbb{Z}[t]/(t^n - 1)$, where $\Gamma_{K,n}$ is the n -cyclic polynomial associated to K and $\Delta_{K,n}$ is the projection of the Alexander polynomial of K on $\mathbb{Z}[t]/(t^n - 1)$.

Proof. To prove the statement we will use the reduced Reidemeister torsion of $M = L(p, q) - N(K)$. In the cellular presentation (2.4) of $\pi_1(M, *)$, the total exponent sum of α in $r(\alpha, \gamma)$ is p , since α is a generator of $\pi_1(L(p, q), *)$. So, by abelianization, we get $H_1(M) = \langle \alpha, \gamma \mid p\alpha + q'\gamma \rangle \cong \mathbb{Z} \oplus \mathbb{Z}_d$, where $d = \gcd(p, q')$. Moreover, if we set $\bar{p} = p/d$ and $\bar{q} = q'/d$ we have that $\eta = \bar{p}\alpha + \bar{q}\gamma$ and $\xi = -s\alpha + r\gamma$ generate, respectively, the torsion part and the free part of $H_1(M)$, where r and s are integers such that $r\bar{p} + s\bar{q} = 1$. Now, consider the covering $F : \bar{M} \rightarrow M$ corresponding to the subgroup $\Phi^{-1}(\text{Tors}(H_1(M)))$ of $\pi_1(M, *)$, where $\Phi : \pi_1(M, *) \rightarrow H_1(M)$ denotes the Hurewitz homomorphism. The Betti number of $H_1(M)$ is one, so the covering F is infinite cyclic with monodromy $\omega_F : H_1(M) \rightarrow \langle t \rangle \cong \mathbb{Z}$ defined by $\omega_F(\xi) = t$ and $\omega_F(\eta) = 1$. Inverting the defining relations for ξ and η we have that $\omega_F(\gamma) = t^{\bar{p}}$ and $\omega_F(\alpha) = t^{-\bar{q}}$. Up to contraction, which does not alter the torsion, the cell complex $C_*(M)$

$$0 \rightarrow C_2 = \langle \varphi(D') \rangle \rightarrow C_1 = \langle \alpha, \gamma \rangle \rightarrow C_0 = \langle * \rangle \rightarrow 0, \quad (4.2)$$

is a cellular decomposition for M . The reduced Reidemeister torsion of M is, up to multiplication by $\pm t^{\pm h}$, the torsion of the complex $\bar{C}_*(\bar{M}) = \mathbb{Q}(t) \otimes_{\mathbb{Z}[t, t^{-1}]} C_*(\bar{M})$,

where $C_*(\overline{M})$ is the lifting of $C_*(M)$, with a fundamental family of cells of $\overline{M}$ as a base. Let $\delta_1 : \overline{C}_1 \rightarrow \overline{C}_0$ and $\delta_2 : \overline{C}_2 \rightarrow \overline{C}_1$ be the boundary operators. If $\tilde{*}$ is the fixed 0-cell over $*$, and $\tilde{\alpha}$ and $\tilde{\gamma}$ are the lifting of, respectively, α and γ with starting point $\tilde{*}$ then, by the action of the monodromy map, we have $\delta_1(\tilde{\alpha}) = (t^{-\bar{q}} - 1)\tilde{*}$ and $\delta_1(\tilde{\gamma}) = (t^{\bar{p}} - 1)\tilde{*}$. Moreover it is well-known (see for example [3, § 9A-9B]) that, the matrix of δ_2 , is the Alexander matrix of the presentation (2.4). So δ_1 and δ_2 are represented by the matrices ${}^t(t^{-\bar{q}} - 1 \quad t^{\bar{p}} - 1)$ and $(Q_\alpha(t) \quad Q_\gamma(t))$, where $Q_\alpha(t)$ (resp. $Q_\gamma(t)$) is obtained from the Fox derivative $\partial r(\alpha, \gamma)/\partial \alpha$ (resp. $\partial r(\alpha, \gamma)/\partial \gamma$) by the substitutions $\alpha = t^{-\bar{q}}$ and $\gamma = t^{\bar{p}}$. The collection $\{1, 1, \emptyset; t^{\bar{p}} - 1, Q_\alpha(t)\}$ is a torsion chain for this complex (see [20, 21]), so $r_M(t) = Q_\alpha(t)/(t^{\bar{p}} - 1) \in \mathbb{Q}(t)$. By Theorem 3.3, $r_M(t) = \Delta_K(t)/(t - 1)$ and so $Q_\alpha(t) = \Delta_K(t)(\sum_{i=0}^{\bar{p}-1} t^i)$, where the equality holds up to units of $\mathbb{Z}[t, t^{-1}]$. To complete the proof we have to show that $Q_\alpha(t) = \Gamma_{K,n}(t^{\bar{p}})$ in $\mathbb{Z}[t]/(t^n - 1)$. Observe that, for each $n > 1$ coprime with p , the monodromy of the unique n -fold strongly-cyclic branched covering of K , F_n , is the composition of ω_F with the epimorphism $\mathbb{Z} \rightarrow \mathbb{Z}_n$, given by $t \rightarrow 1$. Then we have $\omega_{F_n}(\alpha) = -\bar{q}$ and $\omega_{F_n}(\gamma) = \bar{p}$. If $\bar{p} \neq 1$, to calculate $\Gamma_{K,n}$ we have to replace F_n with the equivalent covering with monodromy ω' , such that $\omega'(\gamma) = 1$, and so $\omega'(\alpha) = -\bar{q}\bar{p}^{-1}$, where $\bar{p}^{-1}$ is the inverse of $\bar{p}$ in $\mathbb{Z}_n$. Let $\bar{r}(x, \gamma) = r(x\gamma^{\omega'(\alpha)}, \gamma)$. It is easy to check, using formula (2.5), that $\Gamma_{K,n}(u)$ is equal to the polynomial obtained from $\partial \bar{r}(x, \gamma)/\partial x$ by the substitutions $x = 1$ and $\gamma = u$. Moreover, since $\partial \bar{r}(x, \gamma)/\partial x = \partial r(x\gamma^{\omega'(\alpha)}, \gamma)/\partial x = (\partial r/\partial \alpha)(x\gamma^{\omega'(\alpha)}, \gamma)$ we have that $\Gamma_{K,n}(u)$ is equal to the polynomial obtained from $\partial r/\partial \alpha$ by the substitutions $\alpha = u^{\omega'(\alpha)} = u^{-\bar{q}\bar{p}^{-1}}$, and $\gamma = u$. So, setting $u = t^{\bar{p}}$ in $\Gamma_{K,n}(u)$, we get $Q_\alpha(t)$, which ends the proof. $\square$

Remark 4.1. In the case of $\mathbf{S}^2 \times \mathbf{S}^1$, the previous theorem does not hold, since, as we have already observed, in this case $p = 0$, so for each $n > 1$, $\gcd(n, p) = n \neq 1$. In fact, if we look at the proof, it is easy to see that, since $\bar{p} = p/d = 0$, the monodromy ω_F of the covering that leads to the calculation of the Alexander polynomial sends a meridian of the knot into t^0 . So, the covering whose monodromy is the composition of ω_F with the projection $\mathbb{Z} \rightarrow \mathbb{Z}_n$, defined by $t \rightarrow 1$, is not strongly-cyclic for any value of $n > 1$.

Observe that, when $p = 1$, K is a $(1, 1)$ -knot in $\mathbf{S}^3$, and we get the same statement as in Proposition 2.1. Moreover, if $\text{Tors}(H_1(\mathbb{L}(p, q) - N(K))) \cong \mathbb{Z}_p$, then the projection of the Alexander polynomial on $\mathbb{Z}[t]/(t^n - 1)$ is equal to the n -cyclic polynomial, up to units.

As a corollary of this theorem, we get a generalization of Theorem 3.1.

Corollary 4.1. *Let $K \subset \mathbb{L}(p, q)$ be a $(1, 1)$ -knot, and let $C_n(K)$ be the n -fold strongly-cyclic branched covering of K , with n coprime with p . Denote with d the order of the torsion subgroup of $H_1(\mathbb{L}(p, q) - N(K))$. Then $H_1(C_n(K))$ is finite if and only if no root of $\Delta_{K,n}$ is an n -root of unity. Moreover if d_n denote the order*

of the torsion subgroup of $H_1(C_n(K))$, we have:

$$d_n = \left| \prod_{\zeta^n=1, \Delta_{K,n}(\zeta) \neq 0} \left(\frac{\Delta_{K,n}}{\Phi} \right) (\zeta) \sum_{j=0}^{\frac{n}{d}-1} \zeta^j \right|, \quad (4.3)$$

where Φ is the product of the distinct cyclotomic polynomials Φ_s such that s divides n and Φ_s divides $\Delta_{K,n}$. In particular, if $H_1(C_n(K))$ is finite, we have:

$$\sharp H_1(C_n(K)) = \left| \prod_{\zeta^n=1} \Delta_{K,n}(\zeta) \sum_{j=0}^{\frac{n}{d}-1} \zeta^j \right|. \quad (4.4)$$

Proof. If ω denotes the monodromy of $C_n(K)$, up to equivalence, we can suppose that $\omega(\gamma) = 1$. Consider the equivalent n -fold strongly-cyclic branched covering $C'_n(K)$ with monodromy $\omega' = \mu_{p/d}\omega$, where $\mu_{p/d} : \mathbb{Z}_n \rightarrow \mathbb{Z}_n$ denotes the multiplication by p/d which is invertible in $\mathbb{Z}_n$. Then the isomorphism $H_1(C_n(K)) \rightarrow H_1(C'_n(K))$ is given by $x_i \rightarrow x_j$ where $j \equiv i(p/d) \pmod n$. As previously observed, the circulant matrix whose first row is given by the coefficients of $\Gamma_{K,n}(t)$ is a presentation matrix for $H_1(C_n(K))$ as a $\mathbb{Z}$ -module, so, the circulant matrix B whose first row is given by the coefficients of $G(t) = \Gamma_{K,n}(t^{p/d})$ is a presentation matrix for $H_1(C'_n(K))$ as a $\mathbb{Z}$ -module. Obviously $H_1(C'_n(K))$ is finite if and only if $H_1(C_n(K))$ is finite and the order of the torsion subgroup of $H_1(C'_n(K))$ is d_n . By the theory of circulant matrices (see [2]), there exists a complex unitary matrix F , called Fourier matrix, such that $FBF^* = D = \text{Diag}(G(\zeta_1), G(\zeta_2), \dots, G(\zeta_n))$, where $\zeta_1, \zeta_2, \dots, \zeta_n$ are the n -roots of the unity. So $H_1(C'_n(K))$ is finite if and only if the rank of B is n , and so if and only if $G(\zeta_i) = \Delta_{K,n}(\zeta_i) \sum_{j=0}^{\frac{n}{d}-1} \zeta_i^j \neq 0$ for each $i = 1, \dots, n$. The first statement follows from the fact that, since $\gcd(p, n) = 1$, we have $\gcd(p/d, n) = 1$, and so $q_{p/d}(\zeta_i) \neq 0$, for $i = 1, \dots, n$, where $q_{p/d}(t) = \sum_{j=0}^{\frac{n}{d}-1} t^j$. Moreover, by [19, Theorem 3.3], we have that $d_n = \left| \prod_{\zeta^n=1, G(\zeta) \neq 0} \left(\frac{G}{\Phi} \right) (\zeta) \right|$, where Φ is the product of the distinct cyclotomic polynomials Φ_s such that s divides n and Φ_s divides G . To end the proof, we have only to show that Φ_s is a divisor of $\Delta_{K,n} q_{p/d}$ if and only if it is a divisor of $\Delta_{K,n}$. This follows from the fact that, if s is a divisor of n , we have $\gcd(s, p/d)=1$, so for each ζ primitive s -root of the unity, $q_{p/d}(\zeta) \neq 0$. $\square$

Another straightforward corollary is the following.

Corollary 4.2. *The Alexander polynomial of a $(1,1)$ -knot $K \subset L(p, q)$ satisfies the relation $\Delta_K(1) = \pm d$, where d is the order of the torsion subgroup of $H_1(L(p, q) - N(K))$.*

Proof. It is enough to observe that $\Gamma_{K,n}(1) = \pm p$. $\square$

An interesting question for future study could be whether any symmetric polynomial in $\mathbb{Z}[t, t^{-1}]$ can be realized as the Alexander polynomial of a $(1,1)$ -knot in a lens space.

Example 4.1. Let $K_{p,q}$ be the trivial knot in $L(p, q)$. We have that $\pi_1(L(p, q) - N(K_{p,q}), *) = \langle \alpha, \gamma \mid \alpha^p \rangle$, (for reference see [5]). Then $p = d = \sharp \text{Tors}(H_1(L(p, q) - N(K_{p,q})))$. So, for each $n > p$ with $\gcd(n, p) = 1$, we have $\Gamma_{K_{p,q}, n} = p = \Delta_{K_{p,q}}$.

We end this paper by observing that formula (2.5) can easily be implemented to find $\Gamma_{K,n}$, and so, by Theorem 4.1, to calculate Δ_K , as shown in the following example.

Example 4.2. By results of [7], each $(1,1)$ -knot K can be represented as $K(a, b, c, r)$, where a, b, c, r are nonnegative integer parameters determining the Heegaard diagram of a $(1,1)$ -decomposition of K . For each $m > 2$, let $K_m = K(1, m-2, 0, 1) \subset L(m-2, 1)$. This family of knots is very interesting since, as proved in [14], the m -fold strongly-cyclic branched covering of K_m , with monodromy that sends γ into 1 and α into 0, is the Neuwirth manifold $\mathcal{N}_m$ of type m . These manifolds were introduced by L. Neuwirth in [18], while in [8] it is proved that $\mathcal{N}_m$ is a Seifert manifold of type $(0; -1; (2, 1), \dots, (2, 1))$, with base $\mathbf{S}^2$, Euler number -1 and m exceptional fibers. From the parametric representation of K_m (see [1] for details) we get

$$\pi_1(L(m-2, 1) - N(K_m), *) = \langle \alpha, \gamma \mid (\alpha\gamma)^{m-1}\alpha^{-1}\gamma \rangle, \quad (4.5)$$

and so

$$H_1(L(m-2, 1) - N(K_m)) = \langle \alpha, \gamma \mid (m-2)\alpha + m\gamma \rangle \cong \begin{cases} \mathbb{Z} & \text{if } m \text{ is odd,} \\ \mathbb{Z} \oplus \mathbb{Z}_2 & \text{if } m \text{ is even.} \end{cases}$$

Notice that K_3 is the trefoil knot in $\mathbf{S}^3$.

If we denote with ω the monodromy of the n -fold strongly-cyclic branched covering of K_m , with $\gcd(n, m-2)=1$ and satisfying the condition $\omega(\gamma) = 1$, from the homology relation we get

$$c = \omega(\alpha) = \begin{cases} -m \cdot \overline{m-2}, & \text{if } m \text{ is odd,} \\ -\frac{m}{2} \cdot \overline{\left(\frac{m-2}{2}\right)}, & \text{if } m \text{ is even,} \end{cases}$$

where $\overline{x}$ denote the inverse of x in $\mathbb{Z}_n$. From the substitution $x = \alpha\gamma^{-c}$ in the presentation (4.5), we get the relator $\bar{r}(x, \gamma) = (x\gamma^{c+1})^{m-1}\gamma^{-c}x^{-1}\gamma$. By formula (2.5), $\pi_1(C_n(K_m), \tilde{*}) = G_n(w)$, where

$$w = \left(\prod_{i=0}^{m-2} x_{1+i(1+c)} \right) x_{2+(m-2)(1+c)}^{-1}$$

(subscripts mod n). So, we get:

$$\Gamma_{K_m, n}(t) = \left(\sum_{i=0}^{m-2} t^{i(1+c)} \right) - t^{1+(m-2)(1+c)}.$$

If m is odd then $p/d = m - 2$ and we get

$$\Gamma_{K_m, n}(t^{m-2}) = \left(\sum_{i=0}^{m-2} t^{-2i} \right) - t^{-(m-2)} = \left(\sum_{i=0}^{m-2} t^{2i} \right) - t^{(m-2)} = \Delta_{K_m, n}(t) \sum_{j=0}^{m-3} t^j,$$

(where the second equality holds up to multiplication for $t^{2(m-2)}$). This means that for each $n > 2(m-2)$ coprime with $m-2$, the representative of Δ_{K_m} in $\mathbb{Z}[t]/(t^n - 1)$ does not depend on n . So, for each $m > 2$ odd

$$\Delta_{K_m}(t) = \frac{(\sum_{i=0}^{m-2} t^{2i}) - t^{(m-2)}}{\sum_{j=0}^{m-3} t^j} = \sum_{i=0}^{m-1} (-t)^i.$$

Analogously, if m is even we have $p/d = \frac{m-2}{2}$ and therefore

$$\Gamma_{K_m, n}(t^{(m-2)/2}) = \left(\sum_{i=0}^{m-2} t^{-i} \right) - t^{-(m-2)/2} = \left(\sum_{i=0}^{m-2} t^i \right) - t^{(m-2)/2} = \Delta_{K_m, n}(t) \sum_{j=0}^{(m-4)/2} t^j.$$

By the same considerations, for each $m > 2$ even

$$\Delta_{K_m}(t) = \frac{(\sum_{i=0}^{m-2} t^i) - t^{(m-2)/2}}{\sum_{j=0}^{(m-4)/2} t^j} = t^{m/2} + 1.$$

Acknowledgements

Work performed under the auspices of the G.N.S.A.G.A. of I.N.d.A.M. (Italy) and the University of Bologna, funds for selected research topics. The author would like to thank Michele Mulazzani and the referee for their helpful suggestions.

References

- [1] H. Aydin, I. Gultekin and M. Mulazzani, *Torus knots and Dunwoody manifolds*, Siberian Math. J. **45** (2004) 1-6.
- [2] M. Barnabei and L. B. Montefusco: *Circulant recursive matrices*, Algebraic combinatorics and computer science. A tribute to Gian-Carlo Rota. 111-127, Springer (2001).
- [3] G. Burde and H. Zieschang, *Knots*, Walter de Gruyter, Berlin-New York (1985).
- [4] A. Cattabriga and M. Mulazzani, *Strongly-cyclic branched coverings of (1,1)-knots and cyclic presentations of groups*, Math. Proc. Cambridge Philos. Soc. **135** (2003) 137-146.
- [5] A. Cattabriga and M. Mulazzani, *(1,1)-knots via the mapping class group of the twice punctured torus*, Adv. Geom. **4** (2004) 263-277.
- [6] A. Cattabriga and M. Mulazzani, *All strongly-cyclic branched coverings of (1,1)-knots are Dunwoody manifolds*, J. London Math. Soc. **70** (2004) 512-528.
- [7] A. Cattabriga and M. Mulazzani, *Representations of (1,1)-knots*, to appear in Fundam. Math., II special issue dedicated to the Proceedings of "Knots in Poland" Conference, [ArXiv GT/0501234](#).

- [8] A. Cavicchioli, *Newwirth manifolds and colourings of graphs*, Aequationes Math., **44** (1992) 168-187.
- [9] D. H. Choi and K. H. Ko, *Parametrizations of 1-bridge torus knots*, J. Knot Theory Ramifications **12** (2003) 463-491.
- [10] R. H. Crowell and R. H. Fox, *Introduction to knot theory*, Springer-Verlag, Berlin (1963).
- [11] D.H. Fox *Free differential calculus III. Subgroups*, Ann. Math.(2) **64** (1956) 407-419.
- [12] H. Fujii, *Geometric indices and the Alexander polynomial of a knot*, Proc. Am. Math. Soc. **128** (1996) 2923-2933.
- [13] L. Grasselli and M. Mulazzani, *Genus one 1-bridge knots and Dunwoody manifolds*, Forum Math. **13** (2001) 379-397.
- [14] L. Grasselli and M. Mulazzani, *Seifert manifolds and $(1,1)$ -knots*, preprint (2005), ArXiv GT/0510318.
- [15] T. Kobayashi and O. Saeki, *The Rubinstein-Scharlemann graphic of a 3-manifold as the discriminant set of a stable map*, Pac. J. Math. **195** (2000) 101-156.
- [16] J. Minkus, *The branched cyclic coverings of 2 bridge knots and links*, Mem. Amer. Math. Soc. **255** (1982) 1-68.
- [17] M. Mulazzani, *Cyclic presentations of groups and cyclic branched coverings of $(1,1)$ -knots*, Bull. Korean Math. Soc. **40** (2003) 101-108.
- [18] L. Neuwirth, *An algorithm for the construction of 3-manifolds from 2-complexes*, Proc. Cambridge Philos. Soc. **64** (1968) 603-613.
- [19] D. S. Silver and S. G. Williams, *Torsion numbers of augmented groups with applications to knots and links*, Enseign. Math. **48** (2002) 317-343.
- [20] V.G. Turaev, *The Alexander polynomial of a three-dimensional manifold*, Mat. USSR Sbornik **26** (1975) 313-329.
- [21] V.G. Turaev, *Introduction to Combinatorial Torsions*, Birkhauser, Basel-Boston-Berlin (2001).